

Plexr: The Revolution In Blepharoplasty

Tsioumas G. Sotiris^{1*}, Georgiadis Nikolaos² & Georgiadou Irini³

¹Private Clinic,
Diagnostic Clinic and Aesthetic Surgery Ophthalmology,
Eye surgeon – Oculoplastic,
President of Hellenic Society of Aesthetic Medicine and Non Invasive Surgery,
Greece.

²Private Clinic,
Professor of Ophthalmology
Aristotle University of Thessaloniki,
Greece.

³Private Clinic,
Ph.D. of Aristotle University of Thessaloniki,
Greece.

Accepted 17 December 2014

ABSTRACT

This study was written after three years of implementation of the innovative technique Plexr, where it was found that Plexr is the latest technology for Blepharoplasty and Non Invasive Eye Lid Surgery. The study was performed in 1000 patients, of whom 800 were women and 200 men. Our results showed that the innovative method of Blepharoplasty with Plexr is not short on anything from the classical method of surgery. There is possibility in most patients who fear surgery and the side effects of it to use the method which causes no pain, incisions, surgical sutures and importantly, delivered 100% of the desired result.

Keywords: Plexr, Non surgical blepharoplasty, sublimation, plasma (4th state of matter), no blood , eye lid surgery, eyelid ptosis, wrinkles.

Introduction

Plexr is a cordless micro-surgical hand operated device that transfers concentrated heat to the treated skin tissues. It uses the difference in voltage between the device and the patient's skin. The difference in voltage generates a small electrical arc, similar to a minute lighting. The small lighting causes the sublimation of the fluids contained in superficial part of the skin, without unwanted heat transmission to the adjacent tissues. Additionally it acts on the superficial layer of the skin preserving the lower layers; this will reduce drastically any potential permanent skin damage that could be caused by the misuse of conventional lasers. Place of 'birth' of the medical device is Rome Via del Buer. 'Capture' invention of new energy form was at the State University "Tor Vergata" Rome, where a Professor with thirty years of experience on the electrosurgical and President of the Italian Society of Aesthetic Medicine and Non Invasive Surgery Giorgio Fippi borrowed the natural phenomenon of lightning mode and with the internationally renowned Engineer Giancarlo Millevolte implemented the idea effectively creating a plasma generator inside the machine (plasma = the fourth state of matter) which safely allows the patient who does not wish to have surgery and the doctor who wants to provide high quality service at low cost and equally and in many cases better results, to see 100% of the desired image.

What is plasma?

Plasma is one of the four fundamental states of matter, the others being solid, liquid, and gas. Plasma has properties unlike those of the other states. Plasma can be created by heating a gas or subjecting it to a strong electromagnetic field applied with a laser or microwave generator. This decreases or increases the number of electrons, creating positive or negative charged particles called ions, and is accompanied by the dissociation of molecular bonds, if present.

The presence of a non-negligible number of charge carriers makes plasma electrically conductive so that it responds strongly to electromagnetic fields. Like gas, plasma does not have a definite shape or a definite volume unless enclosed in a container. Unlike gas, under the influence of a magnetic field, it may form structures such as filaments, beams and double layers.

Plasma is the most abundant form of ordinary matter in the Universe, most of which is in the rarefied intergalactic regions, particularly the intracluster medium, and in stars, including the Sun. A common form of plasmas on Earth is seen in neon signs.

Much of the understanding of plasmas has come from the pursuit of controlled nuclear fusion and fusion power, for which plasma physics provides the scientific basis.

Corresponding Author: Tsioumas G. Sotiris^{1*}

Private Clinic, Diagnostic Clinic and Aesthetic Surgery Ophthalmology, Eye surgeon – Oculoplastic, President of Hellenic Society of Aesthetic Medicine and Non Invasive Surgery, Greece..

Email address: s_tsioumas@hotmail.com

Basic vantage point for cases of eyelid surgery, Plexr is a dynamic non-invasive method that allows the operator to urge the patient at any time during the proceedings to open his eyes, highlighting aspects of skin that still need to be treated and avoiding going after gradual and controlled maneuvers that may have cost image lagophthalmos or excessive opening of the eyes. Regarding any corrective surgery the obvious advantage of Plexr is that is also the perfect choice to avoid further suffering of the patient with correction in his private clinic in infinitesimal time. Histological studies from the University of Cheti and the University of Athens (Department of Pathology-Anatomy) also proved that the spots of Plexr not pass the basement membrane of skin.

Purpose of the study

In surgical techniques of upper eyelid with a scalpel, a radio scalpel or laser, the patient must enter a sterilized operating room, to be anesthetized with xylocaine adrenaline injection and this is done by having incisions and sutures.

The purpose of the study is to demonstrate that the method of non-invasive eyelid with Plexr enables the medical treatment of patient without the use of anesthesia (using only anesthetic cream) and sterilization and most importantly,

after treatment the patient leaves without incisions and sutures.

Methodology

The study was conducted over a period of 2 years and used personal medical records. The upper eyelid blepharoplasty was performed to 1000 persons (800 women and 200 men). Target group: 23-82 years old (whether they were smokers or not, had large or small excess skin). We used only a medical device (Plexr). All patients accepted antisepsis, and then photos were taken of their face before the treatment and profile in order to see the difference after the treatment (Picture 1). The next step was to apply the anesthetic cream (Picture 2), wait for 30 minute until the cream act so the patient won't perceive any pain while the treatment is going on. After the action of the cream, we remove it and apply the technique first on the right eyelid (Picture 3) and then on the left. The point where there is excess skin are sublimated. We did not enter wrinkles but areas of excess skin. For the machine to operate, it should not touch the skin as ionization gas are needed, so while the micro spots were pulled out, we leave little spaces of healthy tissue. In each session we sublimated 30% of the excess skin (Picture 4, 5, 6).

Picture 1: Before Plexr



Source: Dr. Tsioumas Sotiris

Picture 2: Before Plexr (With Anesthetic Cream)



Picture 3: During the Application on the Right Eye



Source: Dr. Tsioumas Sotiris

Picture 4: Immediately After Plexr

Source: Dr. Tsioumas Sotiris

Picture 5: (appear crusts)**Picture 6:** After Plexr (No Change in Look)

Source: Dr. Tsioumas Sotiris

Results


800 out of the 1000 persons have achieved 100% of the desired result in three sessions. 200 out of 1000 persons in one or two sessions. Compared with classical blepharoplasty surgical, outcomes were excellent without sutures and

incisions, ectropion and entropion, slanted eyes, lagofthalmos and other complications. The recovery takes place in a shorter period of time (7-15 days) and allows the patients to return to their activities even after treatment. It is important to note that in any of the 1000 cases of patients, there was no complication as referred to above.

Picture 7: Before Plexr

Source: Dr. Tsioumas Sotiris

Picture 7 & 8: After Plexr 2 Sessions



Picture 9: After Plexr 2 Sessions



After Two Years of Plexr Application



After Two Years of Plexr Application



Discussion

The need of the physician to provide improved innovative non invasive techniques to patients with less pain and cost, but

also the need of the patient for cosmetic procedures that avoided them in the past for fear of surgery. The major advantage of Plexr is security that it provides to the patient on undesirable effects. This was proved by the results of the

1000 incidents were subjected to bloodless Blepharoplasty with Plexr. Patients being perfectly pleased with the result, recommend the method to other patients. The Plexr is the bloodless revolution in Blepharoplasty because the doctor can do as many incidents as he wishes per day in his private practice, even take it to a cooperating office or clinic, without side effects and with the best outcome for the patient.

Conclusion

Given the fact that Plexr treatment is something totally new, it is obvious that doctor has to take some more time to explain although the method is well approved by the patients once tried. Last but not least a good percentage of our sample is bound to repeat the procedure in other parts of their face resulting in satisfaction for the patients getting a natural result (their look didn't change) and more profit for the doctor.

References

- M. Ceccarelli Invecchiamento generale e cutaneo in medicina estetica
- Trattato di medicina estetica Professor Alberto Massirone edizioni Piccin.
- Chang YC, Yang SF, Tai KW, Chou MY, Hsieh YS. Increased tissue inhibitor of metalloproteinase-1 expression and inhibition of gelatinase A activity in buccal mucosal fibroblasts by arecoline as possible mechanisms for oral submucous fibrosis.
- Denton CP, Abraham DJ. Transforming growth factor-beta and connective tissue growth factor: key cytokines in scleroderma pathogenesis.
- Jelaska A, Strehlow D, Korn JH Fibrotic and normal fibroblast of type I and type III procollagen mRNA in cultured fibroblasts of patients with incisional hernia.
- Leask A, Holmes A, Abraham DJ. Connective tissue growth factor: a new and important player in the pathogenesis of fibrosis.
- Lesley J, Hascall VC, Tammi M, Hyman R. Hyaluronan binding by cell surface CD44.
- Lu Y, Luo S, Liu J. The influence of transforming growth factor beta 1 (TGF beta 1) on fibroblast proliferation and collagen synthesis
- Sato M, Shegogue D, Gore EA, Smith EA, McDermott PJ, Trojanowska M. Role of p38 MAPK in transforming growth factor beta stimulation of collagen production by scleroderma and healthy dermal fibroblasts.
- Si Z, Rhanjit B, Rosch R, Rene PM, Klosterhalfen B, Klinge U. Impaired balance
- Fisher GJ (2005). "The Pathophysiology of Photoaging of the Skin." *Cutis*, Feb;75(2S):5-9.
- Fisher, G.J., Wang, Z.Q., Datta, S.C. et al (1977). Pathophysiology of premature skin aging induced by ultraviolet light. *N. Engl. J. Med.*; 337(20): 419-29.
- G.J., Wang, Z.Q., Datta, S.C., Varani, J., Kang, S., Voorhees, J.J. (1997) Pathophysiology of premature skin aging induced by ultraviolet light. *N. Engl. J. Med.*; 337(20): 1419-28.
- Hackenbrock C.R., Chazotte B., Gupte S.S. (1986), The random collision model and a critical assessment of diffusion and collision in mitochondrial electron transport. *J. Bioenerg. Biomembr.* 18 331-368.
- Schägger H., Pfeiffer K (2001) . The ratio of oxidative phosphorylation complexes I-V in bovine heart mitochondria and the composition of respiratory chain supercomplexes. *J. Biol. Chem.* 276; 37861-37867.
- Bianchi C., Genova M.L., Parenti Castelli G., Lenaz G (2004). The mitochondrial respiratory chain is partially organized in a supercomplex assembly: kinetic evidence using flux control analysis. *J. Biol. Chem.* 279 36562-36569.
- Genova M.L, Baracca A., Biondi A., Casalena G., Faccioli M., Falasca A.I., Formiggini G., Sgarbi G., Solaini G., Lenaz G (2008) . Is supercomplex organization of the respiratory chain required for optimal electron transfer activity? *Biochim. Biophys. Acta.* 1777(7-8) 740-6.
- Chan D.C. (2006), Mitochondria: dynamic organelles in disease, aging, and development. *Cell* 125 1241-1252.
- Lenaz G., Baracca A., Fato R., Genova M.L., Solaini G (2006). New insights into structure and function of mitochondria and their role in ageing and disease. *Antioxid. Redox Signal.* 8; 417-437.
- Lenaz G. (1998), Role of mitochondria in oxidative stress and ageing. *Biochim. Biophys. Acta* 1366; 53-67.
- McFarland R., Taylor R.W., Turnbull D.M (2007). Mitochondrial disease--its impact, etiology, and Pathology. *Curr. Top. Dev. Biol.* 77; 113-155.
- Reeve A.K., Krishnan K.J., Turnbull D.M (2008). Age related mitochondrial degenerative disorders in humans. *Biotechnol. J.* 3 ;750-756.
- Wallace D.C. (2005), A mitochondrial paradigm of metabolic and degenerative diseases, aging, and cancer: a dawn for evolutionary medicine. *Annu. Rev. Genet.* 39; 359-407.
- Zeviani M., Carelli V (2007) . Mitochondrial disorders. *Curr. Opin. Neurol.* 20; 564-571.
- Wang Q, Lu K, Yang L Effects of hyaluronic acid-stimulating factor on viability and collagen synthesis of fibroblasts.
- Ghersetich I., Comacchi C, Lotti. T (1999). Ringiovanire la pelle: peeling, dermoabrasione, laser. In. *Dermatologia e medicina interna*. Eds. G. Palminteri;, R. Scerrato, T. Lotti, M. Brai. Casa Editrice Mattioli Pavia, pp1349-1355.
- Ghersetich I, Teofoli P, Ribuffo M, Lotti T (1999). Chemical peelings..Eds. A.D. Katsambas, T.M. Lotti Springer, pp 645-656.
- IL PLEXR Medical Device, CE-0434 Devices for Dermatological and Plastic Microsurgery